

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



(19)

(11) Publication number: **61165327 A**

Generated Document.

PATENT ABSTRACTS OF JAPAN(21) Application number: **61006117**(51) Intl. Cl.: **A61K 31/47**(22) Application date: **14.01.86**

(30) Priority:	17.01.85 US 85 692309	(71) Applicant:	mitsubishi chem ind ltd
(43) Date of application publication:	26.07.86	(72) Inventor:	UMETSU KOHEI YUASA SATOSHI NOMURA TATSUO
(84) Designated contracting states:		(74) Representative:	





**(54) REMEDY FOR
HEPATOPATHY**

(57) Abstract:

PURPOSE: To provide the titled remedy for oral administration after meal and containing tritoqualine or its salt as an active component.

CONSTITUTION: A remedy for hepatopathy can be produced by using the tritoqualine or its salt (e.g. inorganic acid salt such as hydrochloric acid or organic acid salt such as acetic acid, etc.) as an active component. The agent is used in the form of tablet, capsule, powder, particle, granule, elixir, etc. A gelatin-type capsule, preferably a light-shielding capsule containing titanium oxide, is used as a solid carrier. The packaging material for the packaging of the powder, particle or granule is preferably a composite film laminated with an aluminum foil. A suspension or syrup containing 0.5W10wt% active component is preferable as a liquid agent. The compound of formula I can be produced by reacting the compound of formula II with the compound of formula III in a solvent, and reducing the resultant compound of formula IV e.g. with tin.

COPYRIGHT: (C)1986,JPO&Japio



還元

I